

32-7947: Recombinant Mouse IL-17A(Discontinued)

Gene : Il17a
Gene ID : 16171
Uniprot ID : Q62386

Description

Source: Human Cells.
MW :16.2kD.

Recombinant Mouse Interleukin-17A is produced by our Mammalian expression system and the target gene encoding Thr22-Ala158 is expressed with a 6His tag at the C-terminus. Interleukin-17 is a potent pro-inflammatory cytokine produced by activated memory T cells. There are at least six members of the IL-17 family in humans and in mice. Mature mouse IL-17A shares 61% and 89% amino acid sequence identity with human and rat IL-17A, respectively. As IL-17 shares properties with IL-1 and TNF-alpha, it may induce joint inflammation and bone and cartilage destruction. This cytokine is found in synovial fluids of patients with rheumatoid arthritis, and produced by rheumatoid arthritis synovium. It increases IL-6 production, induces collagen degradation and decreases collagen synthesis by synovium and cartilage and proteoglycan synthesis in cartilage. IL-17 is also able to increase bone destruction and reduce its formation. Blocking of interleukin-17 with specific inhibitors provides a protective inhibition of cartilage and bone degradation.

Product Info

Amount : 17A / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : NQGSVAGSCSCDRTISSGTQIPQGTLDHIRKYLKAFHRCPPFFIRFQLQSKSVCGGSQDQWVRELVD
CFERKECGTGHGKSFHHQKHLPQASTQTPEAAEGTPSDTSTPAHSQSTQHSSTLPSGALSLNKEH
TQPWEMTTLPSTGYGLEARPEAEANEKQQDDRQQEAPGAGASTPAWVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.