

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7957: Recombinant Mouse Cathepsin L/CTSL (C-6His)(Discontinued)

Gene : Ctsl **Gene ID :** 13039 **Uniprot ID :** P06797

Description

Source: Human Cells.

MW:36.8kD.

Recombinant Mouse Cathepsin L is produced by our Mammalian expression system and the target gene encoding Thr18-Asn334 is expressed with a 6His tag at the C-terminus. Mouse Cathepsin L is a lysosomal cysteine proteinase which is a member of the peptidase C1 family. This protein is a dimer composed of disulfide-linked heavy and light chains, both produced from a single protein precursor. Cathepsin L plays a major role in intracellular protein catabolism. Its substrates include collagen and elastin, as well as alpha-1 protease inhibitor, a major controlling element of neutrophil elastase activity. Cathepsin L has been implicated in several pathologic processes, including myofibril necrosis in myopathies and in myocardial ischemia, and in the renal tubular response to proteinuria.

Product Info

Amount: 6His) / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: TPKFDQTFSAEWHQWKSTHRRLYGTNEEEWRRAIWEKNMRMIQLHNGEYSNGQHGFSMEMNAF

GDMTNEEFRQVVNGYRHQKHKKGRLFQEPLMLKIPKSVDWREKGCVTPVKNQGQCGSCWAFSA SGCLEGQMFLKTGKLISLSEQNLVDCSHAQGNQGCNGGLMDFAFQYIKENGGLDSEESYPYEAKD GSCKYRAEFAVANDTGFVDIPQQEKALMKAVATVGPISVAMDASHPSLQFYSSGIYYEPNCSSKNL DHGVLLVGYGYEGTDSNKNKYWLVKNSWGSEWGMEGYIKIAKDRDNHCGLATAASYPVVNVDHH

HHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.