

32-8020: Recombinant Human Receptor Tyrosine-Protein Kinase ErbB-3/HER3 (C-Fc)(Discontinued)

Gene : ERBB3

Gene ID : 2065

Uniprot ID : P21860

Description

Source: Human Cells.

MW :61.6kD.

Recombinant Human ErbB3 is produced by our Mammalian expression system and the target gene encoding Ser20-Cys331 is expressed with a Fc tag at the C-terminus. Receptor tyrosine-protein kinase erbB-3 is an enzyme that in humans is encoded by the ERBB3 gene. This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. ERBB3 belongs to the protein kinase superfamily, tyr protein kinase family and EGF receptor subfamily. It contains 1 protein kinase domain and it is expressed in Epithelial tissues and brain. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity.

Product Info

Amount : Fc) / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.
Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : SEVGNSQAVCPGTLNGLSVTGDAENQYQTLTKLYERCEVVMGNLEIVLTGHNADLSFLQWIREVT
GYVLVAMNEFSTLPLPNLRVVRGTQVYDGKFAIFVMLNYNTNSSHALRQLRLTQLTEILSGGVYIEK
NDKLCHMDTIDWRDIVRDRDAEIVVKDNGRSCPPCHEVCKGRCWGPSEDQTLTKTICAPQCN
GHCFGNPNQCCHECAGGCSGPQDTCFACRHFNDSGACVPRCPQLVYNKLTFLQLEPNPHT
KYQYGGVVCVASCPhNFVVDQTSVVRACPPDKMEVDKNGLKMCEPCGGLCPKAFVDDIEGRMDE
PKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG
VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQ
VYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVD
KSRWQQGNVVFSCSVMEALHNHYTQKSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.