

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8072: Recombinant Human Ubiquitin-Associated Protein 1/UBAP1/NAG20 (C-6His)

Gene ID: UBAP1
Gene ID: 51271
Uniprot ID: Q9NZ09

Description

Source: E.coli. MW:56.15kD.

Recombinant Human Ubiquitin-Associated Protein 1 is produced by our E.coli expression system and the target gene encoding Met1-Ser502 is expressed with a 6His tag at the C-terminus. Ubiquitin-Associated Protein 1 (UBAP1) belongs to the UBA domain family. Members of this family are related to ubiquitin and the ubiquitination pathway. Because of their cytogenetic location, this UBA domain family member is being studied as a putative target for mutation in nasopharyngeal carcinomas. UBAP1 is highly expressed in the heart, brain, placenta, lung, skeletal muscle, liver, and pancreas. UBAP1 consists of two UBA domains and one UMA domain. The ubiquitin associated domain is throught to be a non-covalent ubiquitin binding domain, including a compact three helix bundle.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 1mM DTT, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MASKKLGADFHGTFSYLDDVPFKTGDKFKTPAKVGLPIGFSLPDCLQVVREVQYDFSLEKKTIEWAEEIKKIEEAE

REAECKIAEAEAKVNSKSGPEGDSKMSFSKTHSTATMPPPINPILASLQHNSILTPTRVSSSATKQKVLSPPHIKA DFNLADFECEEDPFDNLELKTIDEKEELRNILVGTTGPIMAQLLDNNLPRGGSGSVLQDEEVLASLERATLDFKP LHKPNGFITLPQLGNCEKMSLSSKVSLPPIPAVSNIKSLSFPKLDSDDSNQKTAKLASTFHSTSCLRNGTFQNSLK PSTQSSASELNGHHTLGLSALNLDSGTEMPALTSSQMPSLSVLSVCTEESSPPNTGPTVTPPNFSVSQVPNMPS CPQAYSELQMLSPSERQCVETVVNMGYSYECVLRAMKKKGENIEQILDYLFAHGQLCEKGFDPLLVEEALEMH

QCSEEKMMEFLQLMSKFKEMGFELKDIKEVLLLHNNDQDNALEDLMARAGASLEHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.