

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8083: Recombinant Human Toll-Interacting Protein/TOLLIP (C-6His)

Gene ID: 54472 Uniprot ID: Q9H0E2

Description

Source: E.coli. MW :31.3kD.

Recombinant Human Toll-Interacting Protein is produced by our E.coli expression system and the target gene encoding Ala2-Pro274 is expressed with a 6His tag at the C-terminus. Toll-Interacting Protein (TOLLIP) is a member of the tollip family. TOLLIP localizes to the cytoplasm. It contains one C2 domain and one CUE domain. TOLLIP is an inhibitory adaptor protein for Toll-like receptors (TLR). The Toll-like receptors pathway is a part of the immune system that recognize structurally conserved molecular patterns of microbial pathogens, resulting in an inflammatory immune response. TOLLIP constitutes a complex with Tom1 to regulate endosomal transferring of ubiquitinated proteins. TOLLIP can negative regulate Toll-like receptors signaling, which may limit the production of proinflammatory mediators during the process of inflammation and infection.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 5mM EDTA, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

Storage condition : Reconstituted protein solution can be stor samples are stable at -20°C for 3 months.

Amino Acid: ATTVSTQRGPVYIGELPQDFLRITPTQQQRQVQLDAQAAQQLQYGGAVGTVGRLNITVVQAKLAKNYGMTRM

DPYCRLRLGYAVYETPTAHNGAKNPRWNKVIHCTVPPGVDSFYLEIFDERAFSMDDRIAWTHITIPESLRQGKVE DKWYSLSGRQGDDKEGMINLVMSYALLPAAMVMPPOPVVLMPTVYQQGVGYVPITGMPAVCSPGMVPVALPP

AAVNAQPRCSEEDLKAIQDMFPNMDQEVIRSVLEAQRGNKDAAINSLLQMGEEPVEHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.