

## 32-8140: Recombinant Human Tropomyosin $\alpha$ -3 Chain/TPM3

**Gene :** TPM3  
**Gene ID :** 7170  
**Uniprot ID :** P06753

### Description

Source: E. coli.  
MW :29kD.

Recombinant Human Tropomyosin  $\alpha$ -3 is produced by our E.coli expression system and the target gene encoding Met1-Met248 is expressed. Tropomyosin  $\alpha$ -3 Chain (TPM3) is a member of the Tropomyosin family. TPM3 exists as a heterodimer consisting of an  $\alpha$  and a  $\beta$  chain. TPM3 plays a central role in association with the Troponin complex and in the calcium dependent regulation of vertebrate striated muscle contraction. Defects in TPM3 are the cause of thyroid papillary carcinoma. Mutations in the TPM3 gene cause autosomal dominant nemaline myopathy, and oncogenes formed by chromosomal translocations involving this locus are linked with cancer.

### Product Info

**Amount :** 10  $\mu$ g / 50  $\mu$ g  
**Content :** Lyophilized from a 0.2  $\mu$ m filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MAGITTIEAVKRKIQVLQQQADDAEERAERLQREVEGERRAREQAEEVASLNRRRIQLVEEELDRAQERLATALQKLEEAKEKADESERGMKVIENRALKDEEKMELQEIQLEAKHIAEEADRYEEVARKLVIIEGDLERTEERAELAESRCREMDEQIRLMDQNLKCLSAAEKYSQKEDKYEEEIKILTDKLKEAETRAEFAERSVAKLEKTIDDLKLC TKEEHLCTQRMLDQTLLDLNEM

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\mu$ g/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/ $\mu$ g (1 IEU/ $\mu$ g) as determined by LAL test.