

32-8212: Recombinant Human Signal-Regulatory Protein gamma/SIRPG/CD172g (C-Fc)

Gene : SIRPG
Gene ID : 55423
Uniprot ID : Q9P1W8

Description

Source: Human Cells.
MW :63.9kD.

Recombinant Human Signal-Regulatory Protein gamma is produced by our Mammalian expression system and the target gene encoding Glu29-Pro360 is expressed with a Fc tag at the C-terminus. Signal-Regulatory Protein Gamma (SIRPG) is a member of the signal-regulatory protein (SIRP) family and also belongs to the immunoglobulin superfamily. SIRPG is detected in the liver, and at very low levels in the brain, heart, lung, pancreas, kidney, placenta, and skeletal muscle. SIRPG is an immunoglobulin-like cell surface receptor. On binding with CD47, SIRPG mediates cell-cell adhesion. Engagement on T-cells by CD47 on antigen-presenting cells results in enhanced antigen-specific T-cell proliferation and costimulates T-cell activation. SIRPG as receptor-type transmembrane glycoproteins is involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : EEELQMIQPEKLLLVTVGKTATLHCTVTSLLPVGPVLWFRGVGPGRELIYNQKEGHFPRVTTVSDLTRNNMDF
SIRISSITPADVGTYTCVKFRKGSPENVEFKSGPGTEALGAKPSAPVVLGPAARTTPEHTVSFTCESHGFSRDI
TLKWFKNGNELSDFQTNVDPTGQSVAYSIRSTARVVLDPWDVRSQVICEVAHVTLQGDPLRG TANLSEAIRVP
PTLEVTQQPMRVGNQVNVTCQVRKFYPQSLQLTWSSENGNVCQRETASTLTENKDGTYNWTWSWFLVNISDQR
DDVVLTCQVKHGDGQLAVSKRLALEVTVHQDKQSSDATPVDDIEGRMDPEKSCDKTHTCPPCPAPELLGGPSV
FLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDW
LNGKEYCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE
NNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.