

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8418: Recombinant Mouse C-X3-C Motif Chemokine 1/CX3CL1/Fractalkine (C-6His)

Gene ID : 20312 **Uniprot ID :** 035188

Description

Source: Human Cells. MW:34.3kD.

Recombinant Mouse C-X3-C motif chemokine 1 is produced by our Mammalian expression system and the target gene encoding Gln25-Arg337 is expressed with a 6His tag at the C-terminus. Fractalkine(CX3CL1) is a single-pass type I membrane protein and belongs to the intercrine delta family. It consists of an extracellular NH2-terminal domain, a mucin-like stalk, a transmembrane a helix, and a short cytoplasmic tail. CX3CL1 exists in two forms: as a membrane-anchored or as a shed 80-95K glycoprotein. Soluble CX3CL1 is generated by limited proteolysis on the cell surface, and a disintegrin and metallopeptidase 10 (ADAM10) and ADAM17/tumor necrosis factor-a-converting enzyme (ADAM17/TACE) participate in this shedding. It has been suggested that ADAM10 acts in the constitutive shedding, and ADAM17 acts in response to cell activation. The protein may play a role in regulating leukocyte adhesion and migration processes at the endothelium.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

Storage condition : Reconstituted protein solution can be stor samples are stable at -20°C for 3 months.

Amino Acid: QHLGMTKCEIMCDKMTSRIPVALLIRYQLNQESCGKRAIVLETTQHRRFCADPKEKWVQDAMKHLDHQAAALT

KNGGKFEKRVDNVTPGITLATRGLSPSALTKPESATLEDLALELTTISQEARGTMGTSQEPPAAVTGSSLSTSEA QDAGLTAKPQSIGSFEAADISTTVWPSPAVYQSGSSSWAEEKATESPSTTAPSPQVSTTSPSTPEENVGSEGQP PWVQGQDLSPEKSLGSEEINPVHTDNFQERGPGNTVHPSVAPISSEETPSPELVASGSQAPKIEEPIHATADPQK

LSVLITPVPDTQAATRVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ \tilde{A} \parallel \hat{A} μ g (1 IEU/ \tilde{A} \parallel \hat{A} μ g) as determined by LAL test.