

32-8441: Recombinant Human Epidermal Growth Factor Receptor/EGFR/ErbB1/HER1 (C-6His)

Gene : EGFR
Gene ID : 1956
Uniprot ID : P00533

Description

Source: Human Cells.
MW :69.6kD.

Recombinant Human EGFR is produced by our Mammalian expression system and the target gene encoding Leu25-Ser645 is expressed with a 6His tag at the C-terminus. The EGFR subfamily of receptor tyrosine kinases is composed of EGFR, ErbB2, ErbB3 and ErbB4. The EGFR shares 43% - 44% aa sequence identity with the ECD of human EGFR subfamily. All these family members are type I transmembrane glycoproteins with an extracellular ligand binding domain. The extracellular ligand binding domain is containing two cysteine-rich domains separated by a spacer region and a cytoplasmic domain containing a membrane-proximal tyrosine kinase domain. Ligand binding could induce EGFR homodimerization and heterodimerization with ErbB2, resulting in cell signaling, heterodimerization tyrosine phosphorylation and kinase activation. It can bind EGF, amphiregulin, TGF- α , betacellulin, epiregulin, HB-EGF, epigen, and so on. Its signaling regulates multiple biological functions including cell proliferation, differentiation, motility, and apoptosis. EGFR can also be recruited to form heterodimers with the ligand-activated ErbB3 or ErbB4. EGFR is overexpressed in different tumors. Several anti-cancer drugs use EGFR as target.

Product Info

Amount : 10 μ g / 50 μ g
Content : Lyophilized from a 0.2 μ m filtered solution of PBS,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : LEEKKVCQGTSNKLTLQGTFFEDHFLSLQRMFNCEVVLGNLEITYVQRNYDLSFLKTIQEVAGYVLIANTVERIP
LENLQIIRGNMYEYNSYALAVLSNYDANKTGLKELPMRNLQEIHLGAVRFSNNPALCNVESIQWRDIVSSDFLSN
MSMDFQNHGSCQKCDPSCPNGSCWGAGEENCQKLTKIICAQQCSGRCRGKSPSDCCHNQCAAGCTGPRE
SDCLVCRKFRDEATCKDTCPLMLYNPTTYQMDVNPEGKYSFGATCVKKCPRNYVTDHGSCVRACGADSYE
MEEDGVRKCKKCEGPCRKVCNGIGIGEFKDSLSINATNIKHFNCTSIGDLHILPVAFRGDSFTHTPLDPQEL
DILKTVKEITGFLLIQAWPENRTDLHAFENLEIRGRTKQHGFSLAVVSLNITSLGLRSLKEISDGDVVISGNKNLC
YANTINWKKLFGTSGQKTKIISNRGENSCKATGQVCHALCSPEGCWGPEDRDCVSCRNVSRGRECVDKCNLLE
GEPREFVENSECIQCHPECLPQAMNITCTGRGPDNCIQCAHYIDGPHCVKTCPAGVMGENNTLVWKYADAGH
VCHLCHPNCTYGCTGPGLEGCTNGPKIPSVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.