

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8467: Recombinant Mouse Adiponectin/Acrp30/AdipoQ (C-6His, Human Cells)

Gene ID: Adipoq
Gene ID: 11450
Uniprot ID: Q60994

Description

Source: Human Cells. MW :27.6kD.

Recombinant Mouse Adiponectin is produced by our Mammalian expression system and the target gene encoding Glu18-Asn247 is expressed with a 6His tag at the C-terminus. Adiponectin is a secreted protein. It is synthesized exclusively by adipocytes and secreted into plasma. Adiponectin is an important adipokine that is involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Adiponectin Stimulates AMPK phosphorylation and activates in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Adiponectin also antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. It inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. Adiponectin may play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex: LMW, MMW or HMW.

Product Info

Amount: 10 μg / 50 μg

Content: Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: EDDVTTTEELAPALVPPPKGTCAGWMAGIPGHPGHNGTPGRDGRDGTPGEKGEKGDAGLLGPKGETGDVGM

TGAEGPRGFPGTPGRKGEPGEAAYVYRSAFSVGLETRVTVPNVPIRFTKIFYNQQNHYDGSTGKFYCNIPGLYYF SYHITVYMKDVKVSLFKKDKAVLFTYDQYQEKNVDQASGSVLLHLEVGDQVWLQVYGDGDHNGLYADNVND

STFTGFLLYHDTNHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ \tilde{A} \parallel \hat{A} μ g (1 IEU/ \tilde{A} \parallel \hat{A} μ g) as determined by LAL test.