

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8481: Recombinant Mouse Coagulation Factor X/F10 (C-6His)(Discontinued)

Gene ID: F10 **Gene ID:** 14058 **Uniprot ID:** 088947

Description

Source: Human Cells.

MW:55.1kD.

Recombinant Mouse Coagulation factor X is produced by our Mammalian expression system and the target gene encoding Gly21-Asn481 is expressed with a 6His tag at the C-terminus. Mouse coagulation factor X / F10 a member of the peptidase S1 family. The mature F10 is composed mostly of two EGF-like domains, one Gla gamma-carboxy-glutamate domain and one peptidase S1 domain. Factor Xa is a vitamin K-dependent plasma protease that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting. The two chains of F10 are formed from a single-chain precursor by the excision of two Arg residues. A single-chain precursor is initially synthesized in the liver. The light and heavy chains are linked together by disulfide bonds. The light chain contains a Gla and two EGF-like domains. The heavy chain corresponds to the serine protease domain. It can form a heterodimer with SERPINA5.

Product Info

Storage condition:

Amount: 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM MES, 150mM NaCl and 1mM CaCl2 pH7.5..

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: GKGVFINRERANNVLARTRRANSFFEEFKKGNLERECMEEICSYEEVREIFEDDEKTKEYWTKYKDGDQCESSP

CQNQGACRDGIGGYTCTCSEGFEGKNCELFVRKLCRLDNGDCDQFCREEQNSVVCSCASGYFLGNDGKSCIS TAPFPCGKITTGRRKRSVALNTSDSELDLEDALLDEDFLSPTENPIELLNLNETQPERSSDDLVRIVGGRECKDG ECPWQALLINEDNEGFCGGTILNEFYILTAAHCLHQARRFKVRVGDRNTEKEEGNEMVHEVDVVIKHNKFQRD TYDYDIAVLRLKTPITFRMNVAPACLPQKDWAESTLMTQKTGIVSGFGRTHEKGRQSNILKMLEVPYVDRNTCK LSTSFSITONMFCAGYEAKLEDACOGDSGGPHVTRFKNTYYVTGIVSWGEGCARKGKYGIYTKVTTFLKWIDRS

MKARVGPTAETPRTAGPPNVDHHHHHH

Application Note

Endotoxin: Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test.




Fig 1: SDS PAGE analysis of Recombinant Mouse Coagulation Factor X/F10 (C-6His). R lane is for Protein in Reducing conditions, NR is for Non-Reducing conditions.