

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8500: Recombinant Mouse Ephrin-B1/EFNB1 (C-Fc-6His)

Gene ID: 13641 **Uniprot ID:** P52795

Description

Source: Human Cells. MW :49.8kD.

Recombinant Mouse Ephrin-B1 is produced by our Mammalian expression system and the target gene encoding Lys30-Ser229 is expressed with a Fc, 6His tag at the C-terminus. Mouse Ephrin-B1 is a single-pass type I membrane protein which belongs to the ephrin family. It contains an ephrin RBD (ephrin receptor-binding) domain, and expressed in heart, placenta, lung, liver, skeletal muscle, kidney and pancreas. Ephrin-B1 is cell surface transmembrane ligand for Eph receptors, a family of receptor tyrosine kinases which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. It binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. It may play a role in cell adhesion and function in the development or maintenance of the nervous system.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

Storage condition : Reconstituted protein solution can be stor samples are stable at -20°C for 3 months.

Amino Acid: KNLEPVSWSSLNPKFLSGKGLVIYPKIGDKLDIICPRAEAGRPYEYYKLYLVRPEQAAACSTVLDPNVLVTCNKPH

QEIRFTIKFQEFSPNYMGLEFKKYHDYYITSTSNGSLEGLENREGGVCRTRTMKIVMKVGQDPNAVTPEQLTTSR PSKESDNTVKTATQAPGRGSQGDSDGKHETVNQEEKSGPGAGGGGSGDSVDDIEGRMDEPKSCDKTHTCPP CPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV VSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKHH

НННН

Application Note

Endotoxin : Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.