

32-8564: Recombinant Human VEGF Receptor 1/VEGF R1/FLT-1 (C-Fc)

Gene : FLT1
Gene ID : 2321
Uniprot ID : P17948

Description

Source: Human Cells.
MW :109kD.

Recombinant Human Vascular Endothelial Growth Factor Receptor 1 is produced by our Mammalian expression system and the target gene encoding Ser27-Asn756 is expressed with a Fc tag at the C-terminus. Human Vascular endothelial growth factor receptor 1(VEGFR-1, FLT-1) is a member of the the class III subfamily of receptor tyrosine kinases (RTKs) and Tyr protein kinase family and CSF-1/PDGF receptor subfamily. VEGFR-1 is widely expressed in human tissues including normal lung, placenta, liver, kidney, heart and brain tissues. It is specifically expressed in most of the vascular endothelial cells and peripheral blood monocytes. VEGFR-1 contains seven Ig-like C2-type domains and one protein kinase domain. VEGFR-1 is an essential receptor tyrosine kinase and plays an important role in the regulation of VEGF family-mediated vasculogenesis, angiogenesis, and lymphangiogenesis. It is also mediators of neurotrophic activity and regulators of hematopoietic development. VEGFR-1 is a receptor for VEGF, VEGFB and PGF. It has a tyrosine-protein kinase activity. Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFA, VEGFB and PGF. It may play an essential role as a negative regulator of embryonic angiogenesis by inhibiting excessive proliferation of endothelial cells and promote endothelial cell proliferation, survival and angiogenesis in adulthood. Its function in promoting cell proliferation seems to be cell-type specific. VEGFR-1 can also promote PGF-mediated proliferation of endothelial cells, proliferation of some types of cancer cells, but does not promote proliferation of normal fibroblasts (in vitro).

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : SKLKDPELSLKGTQHIMQAGQTLHLQCRGEAAHKWLSPEMVSKESERLSITKSACGRNGKQFCST
LTLNTAQANHTGFYSCKYLAVPTSKKKKETESAIYIFISDTGRPFVEMYSEIPEIIHMTGRELVIPCRV
TSPNITVTLKKFPLDTLIPDGKRIWDSRKGFISNATYKEIGLLTCEATVNGHLYKTNLYLTHRQTNTIID
VQISTPRPVKLLRGHTLVLNCTATTPLNTRVQMTWSYPDEKNKRASVRRRIDQSNSHANIFYSVLTI
DKMQNKDKGLYTCRVRSGPSFKSVNTSVHIYDKAFITVKHRKQQVLETVAGKRSYRLSMKVKAFF
SPEVVWLKDGLPATEKSARYLTRGYSLIKDVTEEDAGNYTILLSIKQSNVFNLTATLIVNVKPKIYE
KAVSSFPDPALYPLGSRQILTCTAYGIPQPTIKWFWHPCNHNHSEARCDFCNNNEESFILDADSNM
GNRIESITQRMAIEGKNKMASTLVVADSRISGIYICIASNKVGTVGRNISFYITDVPNGFHVNLKMP
TEGEDKLSCVTNKFYRDVTWILLRTVNNRTMHYSISKQKMAITKEHSITLNLTIMNVSLQDSGTYA
CRARNVYTGEELQKKEITIRDQEAPYLLRNLS DHTVAISSSTTL DCHANGVPEPQITWFKNNHKKIQQ
EPGILGPSSTLFIERVTEEDGVYHCKATNQKGSVESSAYLTVQGTSDKSNVDDIEGRMDPEKS
CDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEV
HNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYT
LPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSR
WQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Endotoxin : Less than 0.1 ng/Âµg (1 IEU/Âµg) as determined by LAL test.