

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8600: Recombinant Mouse Leukocyte Mono Ig-Like Receptor 1/LMIR1/CD300a (C-Fc)(Discontinued)

Gene: Cd300a Gene ID: 217303 Uniprot ID: Q6SJQ0

Description

Source: Human Cells.

MW:44.3kD.

Recombinant Mouse LMIR1 is produced by our Mammalian expression system and the target gene encoding Leu28-Arg183 is expressed with a Fc tag at the C-terminus. LMIR1, also termed CD300a, is a type I transmembrane glycoprotein with a single IgV-like extracellular domain and an extended membrane proximal region that links the immunoglobulin (Ig) and transmembrane domains and belongs to the immunoglobulin superfamily. The intracellular domain of LMIR1 contains several immunoreceptor tyrosine-based inhibition motifs (ITIMs). When cross-linked, it will be tyrosine phosphorylated and capable of recruiting tyrosine phosphatases (SHP-1, SHP-2) and inositol polyphosphate 5-phosphatase, SHIP. LMIR1 will regulate mast cell-mediated inflammatory responses. LMIR1 is broadly expressed on myeloid and lymphoid cells, and its expression is differentially regulated depending on the cell type.

Product Info

Storage condition:

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: LHGPSTMSGSVGESLSVSCRYEEKFKTKDKYWCRVSLKILCKDIVKTSSSEEARSGRVTIRDHPDN

LTFTVTYESLTLEDADTYMCAVDISLFDGSLGFDKYFKIELSVVPSEDPVSSPGPTLETPVVSTSLPT KGPALGSNTEGHREHDYSQGLRVDDIEGRMDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKD TLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVE WESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLS

PGK

Application Note

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.