

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8656: Recombinant Mouse Exostosin-Like 2/EXTL2 (N-6His)

**Gene ID:** 58193 **Uniprot ID:** Q9ES89

## **Description**

Source: Human Cells.

MW:33.6kD.

Recombinant Mouse Exostosin-like 2 is produced by our Mammalian expression system and the target gene encoding Asn43-Met330 is expressed with a 6His tag at the N-terminus. Exostosin-like 2 (EXTL2) is a member of the exostosin (EXT)-related family which contains five members: EXT1, EXT2, EXTL1, EXTL2, and EXTL3. Studies have shown that EXT gene family members have the activities of heparan sulfate-synthesizing glycosyltransferases. EXT1 and EXT2, which have been identified as causal genes for hereditary multiple exostoses, have HS-GlcAT-II and GlcNAcT-II activities. EXTL1 has GlcNAcT-II activity and EXTL3 has GlcNAcT-I and -II activities. EXTL2 has GlcNAcT-I and N-acetylgalactosaminyltransferase activities, and transfers a GlcNAc residue to the tetrasaccharide linkage region when this region is phosphorylated by a xylose kinase 1 (FAM20B) and thereby terminate chain elongation. In mice, lack of EXTL2 causes glycosaminoglycan (GAG) overproduction and structural changes of GAGs associated with pathological processes.

## **Product Info**

**Amount:** 10 μg / 50 μg

**Content:** Lyophilized from a 0.2 µm filtered solution of 20mM Tris,150mM NaCl,pH8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: HHHHHHHNIKEDKMLTLRREIKSPSKSALDSFTLIMQTYNRTDLLLRLLNHYQAVPSLHKVIVVWNNVGEKGPEEL

WNSLGPHPIPVIFKPQTANKMRNRLQVFPEVETNAVLMVDDDTLISAQDLVFAFSIWQQFPDQIIGFVPRKHVST SSGIYSYGGFELQTPGPGNGDQYSMVLIGASFFNSKYLELFQKQPAAVHALIDETQNCDDIAMNFLVTRHTGKP SGIFVKPINMVNLEKETNGYSGMWHRAEHFLQRSYCINKLVNIYDGMPLKYSNIMISQFGFPYANHKSKM

## **Application Note**

**Endotoxin :** Less than 0.1 ng/ $\tilde{A}$  $\parallel$  $\hat{A}$  $\mu$ g (1 IEU/ $\tilde{A}$  $\parallel$  $\hat{A}$  $\mu$ g) as determined by LAL test.