9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-8666: Recombinant Mouse Serpin D1/Heparin Cofactor II/HCF2 (C-6His)

Gene : Serpind1
Gene ID : 15160
Uniprot ID : P49182

Description

Source: Human Cells.
MW :53.1kD.
Recombinant Mouse Serpin D1/Heparin cofactor 2/HCF2 is produced by our Mammalian expression system and the target gene encoding Glu24-Ser478 is expressed with a 6His tag at the C-terminus. SerpinD1, also known as heparin cofactor II(HC-II), is a member of Serpin superfamily of the serine proteinase inhibitors. It is a single chain glycoprotein with a size of 66.5 kDa and is secreted from hepatocytes. HC-II acts as a thrombin inhibitor in the coagulation cascade, in a glycosaminoglycan-dependent pathway using the release of a sequestered hirudin-like N-terminal tail for interaction with thrombin. This serpin belongs to multiple member group V2 of vertebrate serpin classification. It has been suggested that HC-II is a predictor of decreased atherosclerosis in the elderly and protective against atherosclerosis in mice. HCII can used as a predictive biomarker and therapeutic target for atherosclerosis.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM HEPES,150mM NaCl,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : EQLTNENLTTSFLPANFHKENTVTNDWIPEGEEDEDYLDLEKLLGEDDDYIYIIDAVSPTDSESSAGNILQLFQGK SRIQRLNILNAKFAFNLYRVLKDQATTSDNLFIAPVGISTAMGMISLGLRGETHEEVHSVLHFRDFVNASSKYEVT TIHNLFRKLTHRLFRRNFGYTLRSVNGLYIQKQFPIREDFKAAMREFYFAEAQEANFPDPAFISKANNHILKLTKGL IKEALENIDPATQMLILNCIYFKGTWVNKFPVEMTHNHNFRLNEREVVKVSMMQTKGNFLAANDQELDCDILQL EYVGGISMLIVVPRKLSGMKTLEAQLTPQVVERWQKSMTNRTREVLLPKFKLEKNYNLVEVLKSMGITKLFNKN GNMSGISDQRIAIDLFKHQSTITVNEEGTQAAAVTTVGFMPLSTQVRFTVDRPFLFLVYEHRTSCLLFMGKVTNP AKSVDHHHHHH

Application Note

Endotoxin : Less than 0.1 ng/ÃÂµg (1 IEU/ÃÂµg) as determined by LAL test.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.