

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8683: Recombinant Mouse Nogo-66 Receptor/Reticulon 4 Receptor/NgR/RTN4R (C-Fc)

**Gene ID:** 8tn4r **Gene ID:** 65079 **Uniprot ID:** Q99PI8

## **Description**

Source: Human Cells. MW:72.7kD.

Recombinant Mouse NgR is produced by our Mammalian expression system and the target gene encoding Cys27-Ser447 is expressed with a Fc tag at the C-terminus. Nogo Receptor (NgR) is a glycosylphosphoinositol (GPI)-anchored protein that belongs to the Nogo recptor family. Human NgR is predominantly expressed in neurons and their axons in the central nervous systems. As a receptor for myelin-derived proteins Nogo, myelin-associated glycoprotein (MAG) and myelin oligodendrocyte glycoprotein (OMG), NgR mediates axonal growth inhibition and may play a role in regulating axonal regeneration and plasticity in the adult central nervous system. NgR may be proposed as a potential drug target for treatment of various neurological conditions. Additionally, NgR may play a role in regulating the function of gap junctions.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

**Content:** Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: CPGACVCYNEPKVTTSCPQQGLQAVPTGIPASSQRIFLHGNRISHVPAASFQSCRNLTILWLHSNALARIDAAAF

TGLTLLEQLDLSDNAQLHVVDPTTFHGLGHLHTLHLDRCGLRELGPGLFRGLAALQYLYLQDNNLQALPDNTF RDLGNLTHLFLHGNRIPSVPEHAFRGLHSLDRLLLHQNHVARVHPHAFRDLGRLMTLYLFANNLSMLPAEVLMP LRSLQYLRLNDNPWVCDCRARPLWAWLQKFRGSSSEVPCNLPQRLADRDLKRLAASDLEGCAVASGPFRPIQ TSQLTDEELLSLPKCCQPDAADKASVLEPGRPASAGNALKGRVPPGDTPPGNGSGPRHINDSPFGTLPSSAEPP LTALRPGGSEPPGLPTTGPRRRPGCSRKNRTRSHCRLGQAGSGASGTGDAEGSVDDIEGRMDEPKSCDKTHT CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYP SDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG

Κ

## **Application Note**

**Endotoxin**: Less than 0.1 ng/ $\tilde{A} \sqcap \hat{A} \mu g$  (1 IEU/ $\tilde{A} \sqcap \hat{A} \mu g$ ) as determined by LAL test.