

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8702: Recombinant Human/Mouse/Rat Irisin/FNDC5 (C-6His)

Gene ID: 252995
Uniprot ID: Q8NAU1

## **Description**

Source: Human Cells. MW:13.6kD.

Recombinant Human/Mouse/Rat Irisin is produced by our Mammalian expression system and the target gene encoding Asp32-Glu143 is expressed with a 6His tag at the C-terminus. Fibronectin type III domain-containing protein 5, the precursor of irisin, is a protein that is encoded by the FNDC5 gene. Human Irisin is synthesized as a 212 amino acid (aa) precursor encoding a type 1 transmembrane protein with a 121 aa extracellular domain (ECD), a 21 aa transmembrane domain, and a 39 aa cytoplasmic domain. The ECD of Irisin contains a fibronectin type III domain and multiple glycosylation sites. The ECD is proteolytically cleaved to release the 112 aa soluble Irisin hormone into circulation. Mature human, mouse share 100% sequence identity. Irisin induces expression of peroxisome proliferatoractivated receptor gamma coactivator 1 alpha (PGC1 alpha) and uncoupling protein1(UCP1), mitochondrialassociated metabolic proteins. Irisin induces the transition of white adipose tissue into more metabolically active beige adipose tissue. Irisin also regulates neuronal cell differentiation and neurite outgrowth in the brain and is involved in the differentiation of osteoblasts.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

**Content:** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: DSPSAPVNVTVRHLKANSAVVSWDVLEDEVVIGFAISQQKKDVRMLRFIQEVNTTTRSCALWDLEEDTEYIVHV

QAISIQGQSPASEPVLFKTPREAEKMASKNKDEVTMKEHHHHHH

## **Application Note**

**Endotoxin :** Less than  $0.1 \text{ ng/}\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.