

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8768: Recombinant Human Tumor Necrosis Factor Receptor I/TNFRSF1A/CD120a (C-Fc)

Gene ID: TNFRSF1A
Gene ID: 7132
Uniprot ID: P19438

Description

Source: Human Cells. MW:47.2kD.

Recombinant Human Tumor Necrosis Factor Receptor I is produced by our Mammalian expression system and the target gene encoding Leu30-Thr211 is expressed with a Fc tag at the C-terminus. Tumor necrosis factor receptor superfamily member 1A (TNFRSF1A) is a member of the tumor necrosis factor receptor superfamily. TNFRSF1A is one of the major receptors for the tumor necrosis factor-alpha. It can activate the transcription factor NF-?B, mediate apoptosis, and function as a regulator of inflammation. Antiapoptotic protein BCL2-associated athanogene 4 (BAG4/SODD) and adaptor proteins TRADD and TRAF2 have been shown to interact with this receptor, and thus play regulatory roles in the signal transduction mediated by the receptor. Germline mutations of the extracellular domains of this receptor were found to be associated with the human genetic disorder called tumor necrosis factor associated periodic syndrome (TRAPS) or periodic fever syndrome.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid: LVPHLGDREKRDSVCPQGKYIHPQNNSICCTKCHKGTYLYNDCPGPGQDTDCRECESGSFTASENHLRHCLSC

SKCRKEMGQVEISSCTVDRDTVCGCRKNQYRHYWSENLFQCFNCSLCLNGTVHLSCQEKQNTVCTCHAGFFL RENECVSCSNCKKSLECTKLCLPQIENVKGTEDSGTTIEGRDMDPKSCDKTHTCPPCPAPELLGGPSVFLFPPKP KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT

PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.