

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8829: Recombinant Mouse Signal-Regulatory Protein a-1/SIRPA/CD172a (C-MIgG2a)

Gene ID: Sirpa **Gene ID**: 19261 **Uniprot ID**: Q6P618

Description

Source: Human cells. MW :65.1kD.

Recombinant Mouse Signal-Regulatory Protein alpha 1 is produced by our Mammalian expression system and the target gene encoding Lys32-Asn372 is expressed with a MIgG2a tag at the C-terminus. SIRPa is a type I transmembrane glycoprotein. It contains two Ig-like C1-type domains and one Ig-like V-type domain. Mouse SIRP alpha ECD shares 61%, 75%, 62%, 61%, and 59% as sequence identity with human, rat, equine, bovine, and porcine SIRP alpha, respectively. SIRPa can express in various tissues, mainly on brain and myeloid cells, including macrophages, neutrophils, dendritic and Langerhans cells. It also can detect in neurons, smooth muscle and endothelial cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRPa acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. SIRPa shows adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. SIRPa engagement generally produces a negative regulatory signal; it may mediate negative regulation of phagocytosis, mast cell activation and dendritic cell activation.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: KELKVTQPEKSVSVAAGDSTVLNCTLTSLLPVGPIRWYRGVGPSRLLIYSFAGEYVPRIRNVSDTTK

RNNMDFSIRISNVTPADAGIYYCVKFQKGSSEPDTEIQSGGGTEVYVLAKPSPPEVSGPADRGIPD QKVNFTCKSHGFSPRNITLKWFKDGQELHPLETTVNPSGKNVSYNISSTVRVVLNSMDVNSKVICE VAHITLDRSPLRGIANLSNFIRVSPTVKVTQQSPTSMNQVNLTCRAERFYPEDLQLIWLENGNVSRN DTPKNLTKNTDGTYNYTSLFLVNSSAHREDVVFTCQVKHDQQPAITRNHTVLGFAHSSDQGSMQT FPDNNATHNWNIEGRMDPEPRGPTIKPCPPCKCPAPNLLGGPSVFIFPPKIKDVLMISLSPIVTCVVV DVSEDDPDVQISWFVNNVEVHTAQTQTHREDYNSTLRVVSALPIQHQDWMSGKEFKCKVNNKDL PAPIERTISKPKGSVRAPQVYVLPPPEEEMTKKQVTLTCMVTDFMPEDIYVEWTNNGKTELNYKNT

EPVLDSDGSYFMYSKLRVEKKNWVERNSYSCSVVHEGLHNHHTTKSFSRTPGK

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.