

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8868: Recombinant Human Retinol-Binding Protein 3 (N-6His)

**Gene ID:** 5949 **Uniprot ID:** P10745

## **Description**

Source: E.coli. MW :35.2kD.

Recombinant Human Retinol-binding Protein 3 is produced by our E.coli expression system and the target gene encoding Thr321-Leu630 is expressed fused with a 6His tag at the N-terminus. Retinol-binding proteins (RBP) are a family of proteins with diverse functions. They are carrier proteins that bind retinol. Retinol and retinoic acid play crucial roles in the modulation of gene expression and overall development of an embryo. However, deficit or excess of either one of these substances can cause early embryo mortality or developmental malformations. Regulation of transport and metabolism of retinol necessary for a successful pregnancy is accomplished via RBP. Retinol binding proteins have been identified within the uterus, embryo, and extraembryonic tissue of the bovine, ovine, and porcine, clearly indicating that RBP plays a role in proper retinol exposure to the embryo and successful transport at the maternal-fetal interface.

## **Product Info**

**Amount:**  $10 \mu g / 50 \mu g$ 

**Content:** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

**Storage condition :** Reconstituted protein solution can be stor samples are stable at -20°C for 3 months.

Amino Acid: MNHKVHHHHHHHHTLRSALPGVVHCLQEVLKDYYTLVDRVPTLLQHLASMDFSTVVSEEDLVTKLNAGLQAAS

EDPRLLVRAIGPTETPSWPAPDAAAEDSPGVAPELPEDEAIRQALVDSVFQVSVLPGNVGYLRFDSFADASVLG VLAPYVLRQVWEPLQDTEHLIMDLRHNPGGPSSAVPLLLSYFQGPEAGPVHLFTTYDRRTNITQEHFSHMELPG PRYSTQRGVYLLTSHRTATAAEEFAFLMQSLGWATLVGEITAGNLLHTRTVPLLDTPEGSLALTVPVLTFIDNHG

EAWLGGGVVPDAIVLAEEALDKAQEVL

## **Application Note**

**Endotoxin :** Less than 0.1 ng/ $\tilde{A}$  $\parallel$  $\hat{A}$  $\mu$ g (1 IEU/ $\tilde{A}$  $\parallel$  $\hat{A}$  $\mu$ g) as determined by LAL test.