

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-8894: Recombinant Mouse ICOS Ligand/B7-H2/CD275 (C-6His)

Gene ID: 1000 costs 
Gene ID: 50723 costs 
Uniprot ID: Q9JHJ8

## **Description**

Source: Human Cells.

MW:27.1kD.

Recombinant Mouse ICOS Ligand is produced by our Mammalian expression system and the target gene encoding Glu47-Lys279 is expressed fused with a 6His tag at the C-terminus. Mouse ICOS ligand(B7-H2) is an approximately transmembrane glycoprotein in the B7 family of immune regulatory molecules. B7-H2 is expressed on antigen presenting cells such as B cells, macrophages, monocytes, and dendritic cells. It binds to ICOS on activated T cells, leading to both positive and negative effects on immune responses including its own down-regulation. The B7-H2 interaction with ICOS is costimulatory for T cell proliferation as well as the development of B cells, plasma cells, follicular helper T cells and germinal centers. B7-H2 contributes to the development of allergic asthma by enhancing Th2 biased immune responses, limiting Th17 responses, and promoting eosinophilic infiltration into the lung. Its activation of ICOS on Treg limits pulmonary inflammation and airway hyperresponsiveness, promotes the development of inhalational tolerance, and impairs antitumor immunity. In the thyroid, B7-H2 is up-regulated on thyrocytes during inflammation and promotes their proliferation and production of thryoid hormones.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: ETEVGAMVGSNVVLSCIDPHRRHFNLSGLYVYWQIENPEVSVTYYLPYKSPGINVDSSYKNRGHLS

LDSMKQGNFSLYLKNVTPQDTQEFTCRVFMNTATELVKILEEVVRLRVAANFSTPVISTSDSSNPG QERTYTCMSKNGYPEPNLYWINTTDNSLIDTALQNNTVYLNKLGLYDVISTLRLPWTSRGDVLCCV

**ENVALHQNITSISQAESFTGNNTKNPQETHNNELKHHHHHH** 

## **Application Note**

Endotoxin : Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.