

## 32-8928: Recombinant Mouse Tumor Necrosis Factor Receptor Superfamily Member 5/CD40(C-6His)

**Gene :** Cd40  
**Gene ID :** 21939  
**Uniprot ID :** P27512

### Description

Source: Human cells.

MW :20.2kD.

Recombinant Mouse Tumor Necrosis Factor Receptor Superfamily Member 5 is produced by our Mammalian expression system and the target gene encoding Leu20-Arg193 is expressed with a 6His tag at the C-terminus. CD40 is a Type I Transmembrane Glycoprotein that belongs to the TNF Receptor Superfamily. CD40 is expressed in B cells, follicular dendritic cells, dendritic cells, activated monocytes, macrophages, endothelial cells, vascular smooth muscle cells, and several tumor cell lines. The extracellular domain of CD40 is characterized by Cysteine rich repeat regions. Interaction of CD40 with its ligand (CD40L) leads to aggregation of CD40 molecules, which in turn interact with cytoplasmic components to initiate signaling pathways. Several different TRAF proteins (adaptor proteins) have been identified to serves as mediators of the signal transduction. CD40 plays an essential role in mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** LGQCVTCSDKQYLHDGQCCDLCQPGSRLTSHCTALEKTQCHPCDSGEFSAQWNREIRCHQHRHCEPNQGLR  
VKKEGTAESDTVCTCKEGQHCTSKDCEACAQHTPCIPGFGVMEMATETTTDTVCHPCPVGFFSNQSSLFKCY  
WTSCEDKNLEVLQKGTSTQTNVICGLKSRMRHHHHHH

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.