

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-8948: Recombinant Cynomolgus CD3d / CD3 delta Protein (C-6His)

Gene ID : CD3D **Gene ID :** 102133701 **Uniprot ID :** Q95LI8

Description

Source: Human Cells. MW:10.4kD.

Recombinant Cynomolgus T-cell Surface Glycoprotein CD3 Delta Chain protein is produced by our Mammalian expression system and the target gene encoding Phe22-Ala105 is expressed with a 6His tag at the C-terminus. T-cell surface glycoprotein CD3 delta chain (CD3D) is a single-pass type I membrane protein. CD3D, together with CD3-gamma, CD3-epsilon and CD3-zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T cell receptor-CD3 complex. CD3 chains are present as CD3gammaepsilon, deltaepsilon, and zetazeta dimers in the receptor complex and play critical roles in the antigen receptor assembly, transport to the cell surface, and the receptor-mediated signal transduction. T cell receptor-CD3 complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. This complex is critical for T-cell development and function, and represents one of the most complex transmembrane receptors. The T cell receptor-CD3 complex is unique in having ten cytoplasmic immunoreceptor tyrosine-based activation motifs(ITAMs). CD3D contains 1 ITAM domain and has been shown to interact with CD8A.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: FKIPVEELEDRVFVKCNTSVTWVEGTVGTLLTNNTRLDLGKRILDPRGIYRCNGTDIYKDKESAVQVHYRMCQN

CVELDPATLAHHHHHH

Application Note

Endotoxin : Less than $0.1 \text{ ng/}\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.