

## 32-8977: Recombinant Human LILRB5/CD85c/LIR-8 (C-6His)(Discontinued)

**Gene :** LILRB5  
**Gene ID :** 10990  
**Uniprot ID :** O75023

### Description

Source: Human Cells.  
MW :47.8kD.

Recombinant Human Leukocyte Immunoglobulin-like Receptor Subfamily B Member 5 is produced by our Mammalian expression system and the target gene encoding Gly24-His456 is expressed with a 6His tag at the C-terminus. Human Leukocyte Immunoglobulin-like Receptor Subfamily B Member 5 (LILRB5/CD85c/LIR-8) belongs to a family of transmembrane glycoproteins that negatively regulate immune cell activation. Mature human LIR-8 consists of a 435 amino acid (aa) extracellular domain with four Iglike domains, a 21 aa transmembrane segment, and a 111 aa cytoplasmic domain with two immunoreceptor tyrosine-based inhibitory motifs (ITIM). Alternative splicing of human LIR-8 generates an isoform that lacks the second Ig-like domain. LIR-8 is expressed on NK cells and in the tryptic granules of mast cells. Following cell activation and degranulation, it is present on the mast cell surface. Activated mast cells may also release soluble forms of LIR-8.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** GTLPKPTLWAEPASVIARGKPVTLWCQGPLETEEYRLDKEGLPWARKRQNPLEPGAKAKFHIPSTVYDSAGRYR  
CYYETPAGWSEPSDPLELVATGFYAETLLALPSPVVASGGNVTLQCDTLDGLLTFVLVEEQKLPRTLYSQKLP  
KGPSQALFPVGPVTPSCRWRFRYYYYRKNPQVWSNPDLLEILVPGVSRKPSLLIPQGSVVARGGSLTLQCRS  
DVGVDIFVLYKEGEHDLVQGGSGQPQAGLSQANFTLGPVSRSHGGQYRCYGAHNLSPRWSAPSDPLDILIAGL  
IPDIPALSVQPGPKVASGENVTLLCQSWHQIDTFFLTKEGAHPPLCLKSKYQSYRHOAEFSMPVTSAQGGTY  
RCYSAIRSYPYLLSSPSYPQELVVSGSPGDPSSLSTGPTPTGPEDQPLTPTGLDPQSGLGRHHHHHHH

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.