

32-8984: Recombinant Human Tumor Necrosis Factor Receptor Superfamily Member 4 (C-Fc-Avi)

Gene : TNFRSF4

Gene ID : 7293

Uniprot ID : P43489

Description

Source: Human Cells.

MW :48.6kD.

Recombinant Human TNF Receptor Superfamily Member 4 is produced by our Mammalian expression system and the target gene encoding Leu29-Ala216 is expressed with a Fc & Avi tag at the C-terminus. OX40, also termed CD134 and TNFRSF4, is a T cell co-stimulatory molecule of the TNF receptor superfamily which plays a key role in the survival and homeostasis of effector and memory T cells. OX40 is expressed on CD4+ and CD8+ T cells upon engagement of the TCR by antigen presenting cells along with co-stimulation by CD40-CD40 Ligand and CD28-B7. The interaction between OX40 and OX40 ligand (OX40L) will occur when activated T cells bind to professional antigen-presenting cells (APCs). The T-cell functions, including cytokine production, expansion, and survival, are then enhanced by the OX40 costimulatory signals. OX40 signals are critical for controlling the function and differentiation of Foxp3+ regulatory T cells. OX40-OX40L interaction regulates T-cell tolerance, peripheral T-cell homeostasis, and T-cell-mediated inflammatory diseases.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : LHCVGDTYPSNDRCCHECRPGNGMVSRCRSQNTVCRPCGPGFYNDVVSSKPCKPCTWCNLRSGSERKQLC
TATQDTVCRCRAGTQPLDSYKPGVDCAPCPGPHFSPGDNQACKPWTNCTLAGKHTLQPASNSSDAICEDRDP
PATQPQETQGPPARPITVQTEAWPRTSQGPSTRPVEVPGGRAVAIEGRMDPKSCDKTHTCPPCPAPELLGGP
SVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ
DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNG
QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGKGLNDIFEAQKIE
WHE

Application Note

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.