

32-9012: Recombinant Mouse Interleukin-2 Receptor Subunit Gamma/IL-2 R gamma/CID γ (C-Fc)

Gene : IL2rg
Gene ID : 16186
Uniprot ID : Q3UPA9

Description

Source: Human Cells.

MW :55.1kD.

Recombinant Mouse Interleukin-2 Receptor Subunit Gamma is produced by our Mammalian expression system and the target gene encoding Ser25-Ala263 is expressed with a Fc tag at the C-terminus. The Interleukin-2 receptor gamma chain (IL-2 R gamma , CD132) of the high affinity functional mouse IL-2 receptor complex is a member the hematopoietin receptor family. It is expressed on most lymphocyte (white blood cell) populations, and its gene is found on the X-chromosome of mammals. Common IL2 receptor- gamma Chain is required for IL-2 receptor signaling. Besides IL-2, the Common IL2 receptor- gamma chain has been shown to be a component of the functional receptor complexes for IL-4, IL-7, IL-9 and IL-15. It is a component of multiple cytokine receptors that are essential for lymphocyte development and function. Common IL2 receptor- gamma Chain is also designated the common gamma chain.

Product Info

Amount : 10 μ g / 50 μ g
Content : Lyophilized from a 0.2 μ m filtered solution of PBS, pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : SKVLMSSANEDIKADLILTSTAPEHLSAPTLPLPEVQCFVFNIEYMNCTWNSSEPAQTNLTLHYRYKVSDNNTFQECSHYLFSEITSGCQIQKEDIQLYQTFVVLQDPQKPQRRVQKLNQLNQLVIPRAPENLTLSNLSESQLELRWKSRHIKERCLQYLQYRSNRDRSWTELIVNHEPRFSLPSVDELKRYTFRVRSRYNPICGSSQWQSKWSQPVHWGSHTVEENPSLFALEAVDDIEGRMDEPKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Endotoxin : Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.