

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-9024: Recombinant Human Butyrophilin Subfamily 2 Member A2/BTN2A2 (C-Fc)

Gene ID: BTN2A2
Gene ID: 10385
Uniprot ID: Q8WVV5

Description

Source: Human Cells.

MW:49.7kD.

Recombinant Human Butyrophilin Subfamily 2 Member A2 is produced by our expression system and the target gene encoding Gln33-Val237 is expressed Butyrophilin 2A2 (BTN2A2) is a widely expressed type I transmembrane glycoprotein that functions as a negative regulator of immune responses. Mature human Butyrophilin 2A2 consisits of a 233 amino acid (aa) extracellular domain with two immunoglobulin-like domains, a 21 aa transmembrane segment, and a 237 aa cytoplasmic domain. Alternative splicing generates additional isoforms of human Butyrophilin 2A2 that lack the first, second, or both Iglike domains as well as isoforms with substitutions and deletions in the cytoplasmic region. Within the immune system, Butyrophilin 2A2 is expressed on thymic epithelial cells, naïve B cells, splenic NK cells, dendritic cells, and peritoneal macrophages and is up-regulated with cell activation. Butyrophilin 2A2 inhibits T cell proliferation and activation and enhances the development of FoxP3+ regulatory T cells. Its up-regulation in the hippocampus is associated with schizophrenia.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20 $^{\circ}$ C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: QFTVVGPANPILAMVGENTTLRCHLSPEKNAEDMEVRWFRSQFSPAVFVYKGGRERTEEQMEEY

RGRITFVSKDINRGSVALVIHNVTAQENGIYRCYFQEGRSYDEAILRLVVAGLGSKPLIEIKAQEDGSI WLECISGGWYPEPLTVWRDPYGEVVPALKEVSIADADGLFMVTTAVIIRDKYVRNVSCSVNNTLLG QEKETVIEGRMDPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED PEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD

GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Application Note

Endotoxin : Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.