

## 32-9024: Recombinant Human Butyrophilin Subfamily 2 Member A2/BTN2A2 (C-Fc)

**Gene :** BTN2A2  
**Gene ID :** 10385  
**Uniprot ID :** Q8WVV5

### Description

Source: Human Cells.  
MW :49.7kD.

Recombinant Human Butyrophilin Subfamily 2 Member A2 is produced by our expression system and the target gene encoding Gln33-Val237 is expressed. Butyrophilin 2A2 (BTN2A2) is a widely expressed type I transmembrane glycoprotein that functions as a negative regulator of immune responses. Mature human Butyrophilin 2A2 consists of a 233 amino acid (aa) extracellular domain with two immunoglobulin-like domains, a 21 aa transmembrane segment, and a 237 aa cytoplasmic domain. Alternative splicing generates additional isoforms of human Butyrophilin 2A2 that lack the first, second, or both Iglike domains as well as isoforms with substitutions and deletions in the cytoplasmic region. Within the immune system, Butyrophilin 2A2 is expressed on thymic epithelial cells, naïve B cells, splenic NK cells, dendritic cells, and peritoneal macrophages and is up-regulated with cell activation. Butyrophilin 2A2 inhibits T cell proliferation and activation and enhances the development of FoxP3+ regulatory T cells. Its up-regulation in the hippocampus is associated with schizophrenia.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** QFTVVGPANPILAMVGENTTLRCHLSPEKNAEDMEVRWFRSQFSPAVFVYKGGRRERTEEQMEEYRGRITFVSK  
DINRGSVALVIHNVTAQENGIYRCYFQEGRSYDEAILRLVVAGLGSKPLIEIKAQEDGSIWLECSGGWYPEPLTV  
WRDPYGEVVPALKEVSIADADGLFMVTTAVIIRDKYVRNVSCSVNNTLLGQEKETVIEGRMDPKSCDKTHTCPP  
CPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV  
VSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDI  
AVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

### Application Note

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.