

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-7551: Monoclonal Antibody to IDH1 (Clone: ABM44B3)

Clonality: Monoclonal
Clone Name: ABM44B3
Application: IHC
Reactivity: Human
Gene: IDH1
Gene ID: 3417

 Gene ID :
 3417

 Uniprot ID :
 075874

 Format :
 Purified

 Alternative Name :
 IDH1,PICD

Isotype: Mouse IgG2a Kappa

Immunogen Information: A partial length recombinant IDH1 protein (amino acids 198-414) was used as the immunogen for

this antibody

Description

Isocitrate dehydrogenase 1 (IDH1) is a dimeric cytosolic NADP-dependent dehydrogenase that catalyzes decarboxylation of isocitrate into alpha-ketoglutarate. It belongs to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). It serves a significant role in cytoplasmic NADPH production. It is found in the cytoplasm and peroxisomes. IDH1 is of great importance in cell metabolism and energy conversion. Mutations in IDH1 are found in a somatic mosaic fashion in patients with multiple enchondromas.

Product Info

Amount : 25 μg / 100 μg

Purification: Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition:

Storage condition:

Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

Immunohistochemical analysis: 5 µg/ml




Fig-1: Immunohistochemical analysis of IDH1 in human testis tissue using IDH1 antibody (Clone: ABM44B3) at 5 µg/ml.