

32-12064: Mouse Fibroblast Growth Factor-acidic

Gene : Fgf1

Gene ID : 14164

Uniprot ID : P61148

Alternative Name : Acidic fibroblast growth factor, Heparin-binding growth factor 1, Fgf-1, Fgfa

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 15.9 kDa (141 aa)

Acidic fibroblast growth factor (FGF-acidic), also known as FGF-1, is a potent inducer of DNA synthesis, cell proliferation, and has chemotactic activities. FGF-acidic regulates cardiogenesis through protein kinase C signaling. FGF-acidic also functions as an insulin sensitizer and mediates adipose tissue remodeling. High serum levels of FGF-acidic are associated with type 2 diabetes mellitus (T2DM), suggesting a pathogenic role of FGF-acidic during T2DM.

Product Info

Amount : 50 µg / 100 µg

Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$

Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, 75 mM sodium chloride, pH 7.5
Sterile water at 0.1 mg/mL

Storage condition : Store at -20°C

Amino Acid : MFNLPLGNYK KPKLLYCSNG GHFLRLPDG TVDGTRDRSD QHIQLQLSAE SAGEVYIKGT ETGQYLAMDT
EGLLYGSQTP NEECLFLERL EENHYNTYTS KKHAEKNWFV GLKKNNGSCKR GPRTHYGQKA ILFLPLPVSS D

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialled to compensate for this loss.



