

32-12074: Mouse Flt-3 Ligand

Gene : Flt3lg
Gene ID : 14256
Uniprot ID : P49772
Alternative Name : SL cytokine, Flt3L, Fms-related tyrosine kinase 3 ligand

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 18.6 kDa (163 aa)

Fms-related tyrosine kinase 3 ligand (FLT-3 ligand) is a growth factor that regulates hematopoietic cell proliferation. FLT-3 ligand signalling is transmitted through the fms-related tyrosine kinase 3 (FLT-3) receptor. FLT-3 ligand promotes the long-term expansion and differentiation of pro-B cells in the presence of interleukin 7 (IL-7,) or in combination of IL-7 and interleukin 3 (IL-3).

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MTPDCYFSHS PISSNFKVKF RELTDHLLKD YPVTVAVNQ DEKHCKALWS LFLAQRWIEQ LKTVAGSKMQ
TLLEDVNTEI HFVTSCTFQP LPECLRFVQT NISHLLKDTC TQLLALKPCI GKACQNFSRC LEVQCQPDSS
TLLPPRSPIA LEATELPEPR PRQ

Application Note

Endotoxin: Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.

Biological Activity was determined by OCI-AML5 cell proliferation at ≤ 10 ng/mL; $\geq 1.0 \times 10^5$ units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80Å°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialled to compensate for this loss.



