

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-12080: Human Galectin-3

Gene: LGALS3 Gene ID: 3958 **Uniprot ID:** P17931

35 kDa lectin, Carbohydrate-binding protein 35, Galactose-specific lectin 3, Galactoside-binding **Alternative Name:** 

protein, IgE-binding protein, L-31, Laminin-binding protein, Lectin L-29, Mac-2 antigen, MAC2

## **Description**

Source: Genetically modified E.coli.

Predicted MW:Â Monomer, 26.2 kDa (250 aa)

 Galectin-3 belongs to the lectin family of carbohydrate binding proteins. Galectin-3 is expressed by a wide range of cell types including activated T cells, tumor cells, macrophages, osteoclasts, fibroblasts, and epithelial cells. Galectin-3 has specific binding affinity for beta-galactoside sugar moieties and has functional roles during development, innate immunity, cell apoptosis, and tumor metastasis. Galectin-3 is associated with cancer, heart failure, stroke, and inflammation. The amino acid sequences of human and mouse Galectin-3 proteins share 80% homology.

## **Product Info**

Amount: 50 μg / 100 μg

**Purification:** Reducing and Non-Reducing SDS PAGE at >= 95%

Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium

Content: phosphate, 50 mM sodium chloride, pH 7.5

Sterile water at 0.1 mg/mL

Storage condition: Store at -20°C

Amino Acid: MADNFSLHDA LSGSGNPNPQ GWPGAWGNQP AGAGGYPGAS YPGAYPGQAP PGAYPGQAPP


> GAYPGAPGAY PGAPAPGVYP GPPSGPGAYP SSGOPSATGA YPATGPYGAP AGPLIVPYNL PLPGGVVPRM LITILGTVKP NANRIALDFO RGNDVAFHFN PRFNENNRRV IVCNTKLDNN WGREEROSVF PFESGKPFKI

QVLVEPDHFK VAVNDAHLLQ YNHRVKKLNE ISKLGISGDI DLTSASYTMI

## **Application Note**

**Endotoxin:** Less than 0.1 ng/ $\tilde{A}$  $\parallel \hat{A}$  $\mu g$  (1 IEU/ $\tilde{A}$  $\parallel \hat{A}$  $\mu g$ ) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80̸°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.





9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

