

32-12104: Mouse GRO-alpha (CXCL1)

Gene :	Cxcl1
Gene ID :	14825
Uniprot ID :	P12850
Alternative Name	Gro, Gro1, Mgsa, Scyb1, C-X-C motif chemokine 1, Platelet-derived growth factor-inducible protein KC, Secretory protein N51, KC-T

Description

Source: Genetically modified E.coli.

Predicted MW:Â Monomer, 7.8 kDa (72 aa)


Growth regulated protein alpha (GRO-alpha), also known as CXCL1, is a chemokine that has mitogenic properties and is a neutrophil chemoattractant. GRO- alpha is secreted by macrophages, epithelial cells, neutrophils, and melanomas. GROalpha signals through the CXCR2 chemokine receptor and is important during spinal cord formation, inflammation, angiogenesis, tumorigenesis, and wound healing.

Product Info	
Amount :	20 μg / 100 μg
Purification :	Reducing and Non-Reducing SDS PAGE at >= 95%
Content :	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) Sterile water at 0.1 mg/mL
Storage condition :	Store at -20°C
Amino Acid :	APIANELRCQ CLQTMAGIHL KNIQSLKVLP SGPHCTQTEV IATLKNGREA CLDPEAPLVQ KIVQKMLKGV PK

Application Note


Endotoxin: Less than 0.1 ng/ \tilde{A} $\hat{A}\mu g$ (1 IEU/ \tilde{A} $\hat{A}\mu g$) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80 \tilde{A} ||ŰC and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss. \tilde{A} ||Å \tilde{A} ||Å





9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



For Research Use Only. Not for use in diagnostic/therapeutics procedures.