

## 32-12141: Mouse Interleukin-3

**Gene :** IL3  
**Gene ID :** 16187  
**Uniprot ID :** P01586

**Alternative Name :** Hematopoietic growth factor, Mast cell growth factor, Multipotential colony-stimulating factor, P-cell-stimulating factor, Csfmu, IL-3

### Description

**Source:** Genetically modified E.coli.

**Predicted MW:** Monomer, 15.2 kDa (135 aa)

Interleukin 3 (IL-3) is a cytokine that is produced by activated T cells and mast cells. IL-3 induces the differentiation of hematopoietic stem cells into myeloid precursor cells, such as erythrocyte, megakaryocyte, granulocyte, monocyte, and dendritic cells. IL-3 also functions in the nervous system and is important during the B-1 cell regulation of chronic inflammatory diseases.

### Product Info

**Amount :** 10 µg / 100 µg  
**Purification :** Reducing and Non-Reducing SDS PAGE at  $\geq 95\%$   
Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)  
**Content :** Sterile water at 0.1 mg/mL  
**Storage condition :** Store at  $-20^{\circ}\text{C}$   
**Amino Acid :** MDTHRLTRL NCSSIVKEII GKLPEPELKT DDEGPSLRNK SFRRVNLSKF VESQGEVDPE DRYVIKSNLQ KLNCLPTSA NDSALPGVFI RDLDDFRKKL RFYMVHLNDL ETVLTSRPPQ PASGSVSPNR GTVEC

### Application Note

**Endotoxin:** Less than  $0.1 \text{ ng}/\mu\text{g}$  ( $1 \text{ IEU}/\mu\text{g}$ ) as determined by LAL test.

Biological Activity was determined by NFS-60 cell proliferation at  $\leq 250 \text{ pg/mL}$ ;  $\geq 4.0 \times 10^6 \text{ units/mg}$ . Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at  $-80^{\circ}\text{C}$  and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



