

32-12147: Human Interleukin-5

Gene : IL5
Gene ID : 3567
Uniprot ID : P05113

Alternative Name : B-cell differentiation factor I, Eosinophil differentiation factor, T-cell replacing factor, TRF

Description

Source: Genetically modified E.coli.

Predicted MW: 13.3/26.6 kDa (116/232 aa)

Interleukin 5 (IL-5) is a hematopoietic growth factor that is expressed in type 2 T helper (Th2) cells, mast cells, and eosinophils. IL-5 acts through the IL-5 receptor (IL-5R), stimulates B cell growth, and mediates eosinophil activation. Human and mouse IL-5 show cross-reactivity.

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium glycinate, pH 8.5
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MIPTIPTSA LVKETLALLS THRTLLIANE TLRIPVPVHK NHQLCTEEIF QGIGTLESQT VQGGTVERLF
KNLSLIKYYI DGQKKKCGEE RRRVNQFLDY LQEFLGVMNT EWIIES

Application Note

Endotoxin: Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.

Biological Activity was determined by TF-1 cell proliferation at ≤ 250 pg/mL; $\geq 4.0 \times 10^6$ units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



