

32-12168: Rat Interleukin-10

Gene : IL10
Gene ID : 25325
Uniprot ID : P29456
Alternative Name : Cytokine synthesis inhibitory factor, CSIF

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 18.8 kDa (161 aa)

Interleukin 10 (IL-10) is an anti-inflammatory cytokine produced by macrophages and type 2 T helper (Th2) cells. IL-10 inhibits the production of pro-inflammatory cytokines such as interferon gamma (IFN γ), tumor necrosis factor alpha (TNF α), interleukin 2 (IL-2), interleukin 3 (IL-3), interleukin 4 (IL-4), and granulocyte-macrophage colony-stimulating factor (GM-CSF), made by macrophages and regulatory T cells. IL-10 also suppresses antigen presentation on antigen presenting cells, and enhances the survival, proliferation, and antibody production of B cells. Mouse IL-10 is not active on human cells.

Product Info

Amount : 20 μ g / 100 μ g
Purification : Reducing and Non-Reducing SDS PAGE at \geq 95%
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MSKGHSIRGD NNCTHFPVSQ THMLRELRAA FSQVKTFQK KDQLDNILLT DSSLQDFKGY LGCQALSEMI
KFYLVEVMPQ AENHGPEIKE HLNSLGEKLLK TLWIQLRRCH RFLPCENKSK AVEQVKNDNFN KLQDKGVYKA
MNEFDIFINC IEAYVTLMKM N

Application Note

Endotoxin: Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.

