9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-12170: Human Interleukin-11

Gene :	IL11
Gene ID :	3589
Uniprot ID :	P20809
Alternative Name :	AGIF (Adipogenesis Inhibitory Factor), oprelvekinÂ

Description


Source: Genetically modified E.coli.
Predicted MW:Â Monomer, 19.3 kDa (179 aa)
Interleukin 11 (IL-11) is a member of the gp130 family of cytokines. IL-11 functions to promote hematopoietic stem cell proliferation and megakaryocyte differentiation. In non-hematopoietic cell populations, IL-11 stimulates acute-phase proteins, modulates the development of immunoglobulin-producing B cells, and regulates bone turnover. IL-11 binds the IL-11Ralpha receptor to activate JAK downstream signaling. Human IL-11 shows activity on murine cells.

Product Info

Amount :	10 µg / 100 µg
Purification :	Reducing and Non-Reducing SDS PAGE at >= 95%
Content :	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) Sterile water at 0.1 mg/mL
Storage condition :	Store at -20°C
Amino Acid :	MPGPPPGPPR VSPDPRAELD STVLLTRSLL ADTRQLAAQL RDKFPADGDH NLDSLPTLAM SAGALGALQL PGVLTRLRAD LLSYLRHVQW LRRAGGSSLK TLEPELGTLQ ARLDRLLRRL QLLMSRLALP QPPPDPPAPP LAPPSSAWGG IRAAHAILGG LHLTLDWAVR GLLLLKTRL

Application Note


Endotoxin: Less than 0.1 ng/Ã□Âµg (1 IEU/Ã□Âµg) as determined by LAL test.
Biological Activity was determined by T11 cell proliferation ORÃ□Â TF-1 cell proliferationat <=2.5 ng/mL; >= 4.0 x 10^5 units/mg OR Ã□Â <=Ã□Â 10 ng/mL; >= 1.0 x 10^5 units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80Ã□Â°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.




Human IL-11 Gel
Figure: 1 ug in each lane (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Human IL-11 has a predicted MW of 19.3 kDa.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com





For Research Use Only. Not for use in diagnostic/therapeutics procedures.