

## 32-12173: Mouse Interleukin-11 (AF)

Gene :	11
Gene ID :	16156
Uniprot ID :	P47873
Alternative Name	AGIF (Adipogenesis Inhibitory Factor), oprelvekinÂ

## Description

**Source:** Genetically modified E.coli.

Predicted MW:Â Monomer, 19.3 kDa (179 aa)

Interleukin 11 (IL-11) is a member of the gp130 family of cytokines. IL-11 functions to promote hematopoietic stem cell proliferation and megakaryocyte differentiation. In non-hematopoietic cell populations, IL-11 stimulates acute-phase proteins, modulates the development of immunoglobulin-producing B cells, and regulates bone turnover. IL-11 binds the IL-11Ralpha receptor to activate JAK downstream signaling.

## **Product Info**

Amount :	10 μg / 100 μg
Purification :	Reducing and Non-Reducing SDS PAGE at $>= 95\%$
Content :	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) Sterile water at 0.1 mg/mL
Storage condition :	Store at -20°C
Amino Acid :	MPGPPAGSPR VSSDPRADLD SAVLLTRSLL ADTRQLAAQM RDKFPADGDH SLDSLPTLAM SAGTLGSLQL PGVLTRLRVD LMSYLRHVQW LRRAGGPSLK TLEPELGALQ ARLERLLRRL QLLMSRLALP QAAPDQPVIP LGPPASAWGS IRAAHAILGG LHLTLDWAVR GLLLLKTRL

## **Application Note**

**Endotoxin:** Less than 0.1 ng/ $\tilde{A}$   $\beta_{\mu}g$  (1 IEU/ $\tilde{A}$   $\hat{A}\mu g$ ) as determined by LAL test.

Biological Activity was determined by T11 cell proliferation at <=2.5 ng/mL; >= 4.0 x 10^5 units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80Å[]ŰC and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.

Reduced:	-	+			
	Contractory	for contractory	MW		
	-		97		
	1		66		
			55		
	100		36		
			31		
			21		
	*		14		
			6		
		Service Statements			
Mouse IL-11 Gel					
Figure: 1 ug in each lane (-) non-					
reducing conditions and (+) reducing					
conditions in a 4-20% Tris-Glycine gel,					
stained with Coomassie Blue. Mouse					
IL-11 has a predicted MW of 19.3 kDa.					



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com

