9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-12176: Mouse Interleukin-13

Gene : Il13
Gene ID : 16163
Uniprot ID : P20109
Alternative Name : T-cell activation protein P600, NC30

Description


Source: Genetically modified E.coli.
Predicted MW:Â Monomer, 12.3 kDa (111 aa)
Interleukin 13 (IL-13) is a cytokine secreted from type 2 T helper (Th2) cells. Â IL-13 has overlapping functions with interleukin 4 (IL-4), including the induction of immunoglobulin E (IgE) secretion from B cells, and the inhibition of interleukin 1 beta (IL-1beta), tumor necrosis factor alpha (TNF-alpha), interleukin 8 (IL-8), and interleukin 6 (IL-6) inflammatory cytokine expression. IL-13 also regulates immune cell inflammation in response to the pathophysiological changes of surrounding non-immune cells. The IL-13 receptor consists of the IL-4Ra and IL-13Ra1 subunits. IL-13 can also bind the IL-13Ra2 receptor with high affinity. IL-13 functions are mediated through the JAK/STAT signaling pathway. Human and mouse IL-13 are cross-reactive. Â

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at >= 95%
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
Sterile 20 mM HCl at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MPVPRSVSLP LTLKELIEEL SNITQDQTPL CNGSMVWSVD LAAGGFCVAL DSLTNISNCN AIYRTQRILH GLCNRKAPTT VSSLPDTKIE VAHFITKLLS YTKQLFRHGP F

Application Note


Endotoxin: Less than 0.1 ng/Ã‚Âµg (1 IEU/Ã‚Âµg) as determined by LAL test.
Biological Activity was determined by TF-1 cell proliferation at <=10 ng/mL; >= 1.0 x 10^5 units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80Ã‚Â°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.



Mouse IL-13
Figure: 1 ug in each lane (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Mouse IL-13 has a predicted MW of 12.3 kDa.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com



For Research Use Only. Not for use in diagnostic/therapeutics procedures.