

32-12183: Mouse Interleukin-17A

Gene :	ll17a
Gene ID :	16171
Uniprot ID :	Q62386
Alternative Name	Ctla8, II17, Cytotoxic T-lymphocyte-associated antigen 8

Description

Source: Genetically modified E.coli.

Predicted MW:Â Dimer, 15.1/30.2 kDa (134/268 aa)

Interleukin 17A (IL-17A), also known as CTLA-8, is a member of the IL-17 family of proteins. IL-17A is a proinflammatory cytokine that is secreted by activated CD4+ and CD8+ T lymphocytes. IL-17A acts through its receptor, IL-17R, to promote increased cytokine and chemokine secretion. In turn, the cytokines and chemokines mediate the immunoregulatory function of IL-17A by promoting the proliferation, maturation, and chemoattraction of neutrophils to inflammatory sites. Elevated levels of IL-17A are associated with rheumatoid arthritis, airway inflammation, allograft rejection, inflammatory bowel disease, psoriasis, cancer, and multiple sclerosis. Human, mouse, and rat IL-17A show activity on mouse cells. Â

Product Info

Amount :	25 µg / 100 µg	
Purification :	urification : Reducing and Non-Reducing SDS PAGE at >= 95%	
Content :	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) Sterile water at 0.1 mg/mL	
Storage condition :	Store at -20°C	
Amino Acid :	MAAIIPQSSA CPNTEAKDFL QNVKVNLKVF NSLGAKVSSR RPSDYLNRST SPWTLHRNED PDRYPSVIWE AQCRHQRCVN AEGKLDHHMN SVLIQQEILV LKREPESCPF TFRVEKMLVG VGCTCVASIV RQAA	

Application Note

Endotoxin: Less than 0.1 ng/ \tilde{A} $\hat{A}\mu g$ (1 IEU/ \tilde{A} $\hat{A}\mu g$) as determined by LAL test.

Biological Activity was determined by Production of IL-6 by mouse 3T3 cells at <= 5 ng/mL; >= 2.0 x 10^5 units/mg. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80Å Δ °C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.

Reduced: +	-
neduced.	- MW
	97
	66
	55
	36
	21
	14
	. 6
Mouse IL-17A	
Figure: 1 up	in each lane (-)
	conditions and (+)
	ditions in a 4-20%
Tris-Glycine	gel, stained with
Coomassie Bl	ue. Mouse IL-17A
is a home	odimer with a
predicted tota	al MW of 30.2kDa.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com



For Research Use Only. Not for use in diagnostic/therapeutics procedures.