

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-12217: Guinea Pig IP-10 (CXCL10)

Gene ID: CXCL10
Gene ID: 100714889
Uniprot ID: H0UXM5

## **Description**

**Source:** Genetically modified E.coli.

**Predicted MW:** Monomer, 8.7 kDa (76 aa)

Interferon gamma-induced protein 10 (IP-10), or CXCL10, is a chemokine secreted by monocytes, endothelial cells and fibroblasts in response to interferon gamma (IFN-gamma). IP-10 functions as a chemoattractant for activated T cells, monocytes, dendritic, and natural killer (NK) cells that express the G protein-coupled receptor CXCR3. IP-10 is an important factor in autoimmune diseases such as Hashimoto's thyroiditis, Graves' disease, and Type 1 diabetes mellitus.

## **Product Info**

**Amount:** 25 μg / 100 μg

**Purification:** Reducing and Non-Reducing SDS PAGE at >= 95%

Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic

**Content:** Acid (TFA)

Sterile water at 0.1 mg/mL

**Storage condition :** Store at -20°C

Amino Acid: IPHSRTIRCT CIETSTQPVN PKSFKKLEII PASQSCPRVE IIATMKMNGE KRCLDPESKV IKNLLKAVRK

**ERSKRS** 

## **Application Note**

**Endotoxin:** Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80 $\mbox{\sc A}$  $\mbox{\sc A}$  $\mbox{\sc A}$  $\mbox{\sc A}$  $\mbox{\sc A}$  $\mbox{\sc A}$  $\mbox{\sc A}$ 

