

32-12229: Mouse Monocyte Chemotactic Protein-1 (CCL2)

Gene : Ccl2
Gene ID : 20296
Uniprot ID : Q5SVU3
Alternative Name : C-C motif chemokine

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 13.8 kDa (125 aa)

Monocyte chemotactic protein 1 (MCP-1), also known as CCL2, is produced by injured or infected tissues. MCP-1 signals through the CCR2 and CCR4 G protein-coupled receptors to recruit memory T cells, monocytes, and dendritic cells to sites of inflammation.

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : QPDAVNAPLT CCYSFTSKMI PMSRLESYKR ITSSRCPEA VVFVTKLKRE VCADPKKEWV QTYIKNLDRN QMRSEPTTLF KTASALRSSA PLNVKLTRKS EANASTTFST TTSSTSVGVT SVTVN

Application Note

Endotoxin: Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.

Biological Activity was determined by THP-1 chemotaxis at ≤ 100 ng/mL. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.



