

32-12232: Mouse Monocyte Chemotactic Protein-2 (CCL8)

Gene : Ccl8
Gene ID : 20307
Uniprot ID : Q9Z121
Alternative Name : C-C motif chemokine 8, Monocyte chemoattractant protein 2, Monocyte chemotactic protein 2, Small-inducible cytokine A8

Description

Source: Genetically modified E.coli.

Predicted MW: Monomer, 8.5 kDa (74 aa)

Monocyte chemotactic protein 2 (MCP-2), also known as CCL8, is a cytokine that is important during allergic and inflammatory responses. MCP-2 activates mast cells, eosinophils, and basophils through the G protein-coupled chemokine receptors CCR1, CCR2B, and CCR5.

Product Info

Amount : 20 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at >= 95%
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : GPDKAPVTCC FHVLLKLIPL RVLKSYERIN NIQCPMEAVV FQTKQGMSLC VDPTQKWVSE YMEILDQKSQLQ

Application Note

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.

Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vial to compensate for this loss.

