

32-12336: Human Vascular Endothelial Growth Factor-165 (AF)

Gene : VEGFA
Gene ID : 7422
Uniprot ID : P15692-4
Alternative Name : Vascular endothelial growth factor A, Vascular permeability factor, VPF

Description

Source: Genetically modified E.coli.

Predicted MW: Dimer, 19.3/38.6 kDa (166/332 aa)

Vascular endothelial growth factor-A (VEGF-A) is produced by a wide variety of cell types, including tumor and vascular cells. VEGF-A is a mediator of vascular growth, vascular permeability, and plays a role in stimulating vasodilation via nitric oxide-dependent pathways. VEGF-A has several alternatively spliced isoforms, with VEGF-165 being the most abundant. The VEGF-165 isoform is a secreted protein that acts on receptors VEGFR-1 and VEGFR-2 to modulate endothelial cell proliferation and angiogenesis.

Product Info

Amount : 10 µg / 100 µg
Purification : Reducing and Non-Reducing SDS PAGE at $\geq 95\%$
Content : Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
Sterile water at 0.1 mg/mL
Storage condition : Store at -20°C
Amino Acid : MAPMAEGGGQ NHHEVVKFMD VYQRSYCHPI ETLVDIFQEY PDEIEYIFKP SCVPLMRCGG
CCNDEGLECV PTEESNITMQ IMRIKPHQGG HIGEMSFLQH NKCECRPKKD RARQENPCGP
CSERRKHLFV QDPQTCKCSC KNTDSRCKAR QLELNERTCR CDKPRR

Application Note

Endotoxin: Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.

Biological Activity was determined by HUVEC Proliferation at ≤ 10 ng/mL; $\geq 1.0 \times 10^5$ units/mg (typical ED50 is < 5 ng/mL). Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialled to compensate for this loss.



