9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-12338: Rat Vascular Endothelial Growth Factor-165

Gene :	Vegfa
Gene ID :	83785
Uniprot ID :	P16612-2
Alternative Name :	Vascular endothelial growth factor A, Â VEGF-A, Vascular permeability factor, VPFÂ Â Â Â

Description


Source: Genetically modified E.coli.
Predicted MW:Â Dimer, 19.4/38.8 kDa (165/330 aa)
Vascular endothelial growth factor-A (VEGF-A) is produced by a wide variety of cell types, including tumor and vascular cells. VEGF-A is a mediator of vascular growth, vascular permeability, and plays a role in stimulating vasodilation via nitric oxide-dependent pathways. VEGF-A has several alternatively spliced isoforms, with VEGF-165 being the most abundant. The VEGF-165 isoform is a secreted protein that acts on receptors VEGFR-1 and VEGFR-2 to modulate endothelial cell proliferation and angiogenesis.

Product Info

Amount :	10 µg / 100 µg
Purification :	Reducing and Non-Reducing SDS PAGE at >= 95%
Content :	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5 Sterile water at 0.1 mg/mL
Storage condition :	Store at -20°C
Amino Acid :	MAPTTEGEQK AHEVVKFMDV YQRSYCRPIE TLVDIFQEYP DEIEYIFKPS CVPLMRCAGC CNDEALECVP TSESNVTMQI MRIKPHQSQH IGEMSFLQHS RCECRPKKDR TKPEKHCEPC SERRKHLFVQ DPQTCKCSCK NTDSRCKARQ LELNERTCRC DKPRR

Application Note


Endotoxin: Less than 0.1 ng/Ã□Âµg (1 IEU/Ã□Âµg) as determined by LAL test.
Biological Activity was determined by Bioactive protein. Centrifuge vial before opening, Suspend the product by gently pipetting the above recommended solution down the sides of the vial. DO NOT VORTEX. Allow several minutes for complete reconstitution. For prolonged storage, dilute to working aliquots in a 0.1% BSA solution, store at -80Ã□Â°C and avoid repeat freeze thaws. Upon reconstitution, a small amount of visible precipitate can be expected. A 10% overfill has been added to the total material vialed to compensate for this loss.



Rat VEGF-165 Gel
Figure: 1 ug run under (-) non-reducing conditions and (+) reducing conditions in a 4-20% Tris-Glycine gel, stained with Coomassie Blue. Rat VEGF-165 is a dimer with a predicted MW of 38.8 kDa (each monomer is 19.4 kDa).

For Research Use Only. Not for use in diagnostic/therapeutics procedures.

9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com



For Research Use Only. Not for use in diagnostic/therapeutics procedures.