

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

10-3014: Monoclonal antibody to TLR5 (Clone: ABM22C2)

Clonality: Monoclonal **Clone Name:** ABM22C2 Application: FACS Reactivity: Human Gene: TLR5 Gene ID: 7100 **Uniprot ID:** 060602 **Purified** Format:

Alternative Name: Toll/interleukin-1 receptor-like protein 3, TIL3

Isotype: Mouse IgG2a, Kappa

Immunogen Information: A partial length recombinant TLR5 protein was used as the immunogen for this antibody.

Description

Toll-like receptor 5 (TLR5) plays a fundamental role in pathogen recognition and downstream signaling to induce effective immunity. It is well expressed in major old tissues including lung, skin, intestine, and spleen. TLR5 is present in synovial of patients with rheumatoid arthritis (RA), act as novel mediator of innate immunity-induced osteoclastogenesis and bone loss. TLR5 is a receptor for flagellin and is present on the basolateral surface of intestinal epithelial cells. It is involved in the migration of intestinal epithelial cells through activation of the p38 MAPK pathway. TLR5-mediated signaling specifically enhanced the maturation of lung dendritic cells. TLR5 influences the progression and outcome of ovarian, sarcoma, and luminal breast tumors. Mutations in TLR5 have been associated with both resistance and susceptibility to systemic lupus erythematosus, and susceptibility to Legionnaire disease.

Product Info

Amount : $25 \mu g / 100 \mu g$

Purification: Protein G Chromatography

Content: 25 μg in 50 μl/100 μg in 200 μl PBS containing 0.05% BSA and 0.05% sodium azide. Sodium

azide is highly toxic.

Storage condition : Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid

repeated freeze and thaw cycles.

Application Note

FlowCytometry: 0.5-1 µg/10^6 Cells



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982 Email: info@abeomics.com




Figure-1: Cell surface flowcytometry analysis of hTLR5 on human PBMC (Monocytes) using 0.5 μ g/10^6 Cells of Anti-TLR5 antibody (Clone: ABM22C2). Green represent isotype control and red represent Anti human TLR5 antibody (Abeomics 10-3014). Goat anti mouse PE conjugated was used as the secondary antibody.