

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

45-1022: Rabbit Polyclonal Antibody to RFP-tag

Clonality: Polyclonal
Application: ELISA
Format: Purified
Isotype: Rabbit IgG

Immunogen Information: RFP epitope tag peptide VNGHEFEIEGEGEGR conjugated to KLH

Description

Well-characterizated antibodies against short-sequence epitope tags are common in the study of protein expression in systems. RFP-tags are composed of 15-residue peptides, VNGHEFEIEGEGEGR, derived from amino acids 22-36 of red fluorescent protein (RFP) from the Discosoma sea anemone.

Product Info

Amount: 40 µg

Purification: Immunoaffinity chromatography

Content: 0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide

The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody

Storage condition: can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below.

Avoid repeated freezing and thawing cycles.

Application Note

ELISA: 0.05-0.2 μg/ml Western blot: 1-2 μg/ml

Reconstitute the lyophilized powder with deionized water (or equivalent) to an antibody concentration of 0.5 mg/ml.




Figure-1 : Western blot analysis of RFP-tag Antibody at 1 μ g/ml on RFP-tagged fusion protein (200 & 40 ng) expressed in E. coli cell lysate.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-2 : Western blot analysis of RFP-tag Antibody at 1 μ g/ml on RFP-tagged fusion protein (40 ng, 8 ng, 1.6 ng, 0.8 ng) expressed in E. coli cell lysate.




Figure-3 : Western blot analysis of RFP-tag Antibody at 1 μ g/ml on red fluorescent protein variants (mCherry, DsRed, DeRed2, mOrange, tdTomato), IRDye 800 Conjugated Goat Anti-Rabbit IgG was used as Secondary Antibody.