

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 45-1034: Rabbit Polyclonal Antibody to Histone H2A(Discontinued)

Clonality: Polyclonal
Application: ELISA
Reactivity: Human
Gene: HIST2H2AC
Gene ID: 8338

 Gene ID:
 8338

 Uniprot ID:
 Q16777

 Format:
 Purified

Alternative Name: Histone 2A, Histone H2A-GL101, Histone H2A/q, H2AFQ

**Isotype:** Rabbit IgG

Immunogen Information: Synthetic peptide derived from the carboxyl terminal of human histone H2A

## **Description**

Histone H2A is the core component of the nucleosome. It is involved in nucleosome wrapping and compacting of DNA into chromatin, limiting DNA's accessibility to the cellular machinery. Histones thereby play a central role in transcription regulation, DNA repair, replication, and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, called histone code, and nucleosome remodeling. Rabbit Anti-Histone H2A Polyclonal Antibody is developed in rabbit using a KLH-coupled synthetic peptide corresponding to the residues near carboxyl terminal of human histone H2A protein (Swiss Prot: Q16777).

## **Product Info**

Amount:  $40 \mu g$ 

**Purification:** Immunoaffinity chromatography

Content: 0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide

The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody

Storage condition: can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or below.

Avoid repeated freezing and thawing cycles.

## **Application Note**

ELISA: 0.05-0.2 μg/ml Western blot: 1-2 μg/ml

Reconstitute the lyophilized powder with deionized water (or equivalent) to an antibody concentration of 0.5 mg/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-1 : Western blot analysis of Histone H2A Antibody at 1  $\mu$ g/ml on Hela cell lysate, 1: Histone H2A Antibody, 2: Histone H2A Antibody pre-incubated with immunizing peptide. IRDye 800 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody.