

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 45-1084: Biotinylated Mouse Monoclonal Antibody to Pembrolizumab (Clone: 10E12C6)

Clone Name: 10E12C6
Application: ELISA
Conjugate: Biotin
Format: Purified

**Isotype :** Mouse IgG1,Kappa **Immunogen Information :** Pembrolizumab

## **Description**

Pembrolizumab (Keytruda) is a humanized monoclonal antibody that is approved by the U.S. Food and Drug Administration for the treatment of patients with unresectable or metastatic melanoma. Pembrolizumab binds to the human cell surface receptor PD-1 (Programmed Cell Death Protein 1) and blocks its interaction with PD-L1, resulting in the activation of T-cell-mediated immune responses against tumor cells. Anti-Pembrolizumab Antibody (10E12C6)[BIOTIN], mAb, Mouse is produced from a hybridoma resulting from the fusion of SP2/0 myeloma and B-lymphocytes obtained from a mouse immunized with Pembrolizumab.

## **Product Info**

**Amount :** 40 μg

**Purification:** Protein A chromatography

**Content :** 0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide.

The antibody is stable in lyophilized form if stored at -20°C or below. The reconstituted antibody

**Storage condition:** can be stored for 2-3 weeks at 2-8°C. For long term storage, aliquot and store at -20°C or

below. Avoid repeated freezing and thawing cycles.

## **Application Note**

ELISA detection: 0.01-0.1  $\mu$ g/ml ELISA blocking: 10-50  $\mu$ g/ml




Figure-1 : ELISA binding of Pembrolizumab antibody (Clone: 10E12C6) binds with Pembrolizumab, Coating antigen: Pembrolizumab at 1  $\mu$ g/ml, Pembrolizumab antibody dilution start from 1,000 ng/ml, EC50= 5.77 ng/ml.



9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com




Figure-2 : ELISA blocking of Pembrolizumab antibody (Clone: 10E12C6) with Pembrolizumab binding with Human PD-1 recombinant protein, Coating antigen: Pembrolizumab at 1  $\mu$ g/ml, PD-1-Fc-biotin final concentration: 20 ng/ml, Anti-Pembrolizumab antibody dilution start from 30  $\mu$ g/ml,IC50= 0.37  $1\frac{1}{4}$ g/ml